

## **1 NAME OF THE MEDICINAL PRODUCT**

Etoricoxib 30 mg film-coated tablets

## **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each film-coated tablet contains 30 mg of etoricoxib.

Excipients with known effect: 1.40 mg lactose monohydrate

For the full list of excipients, see section 6.1.

## **3 PHARMACEUTICAL FORM**

Film-coated tablet.

Blue-green, apple-shaped, biconvex film coated tablets engraved '30' on one face and other face plain, approximately 6.4 mm length and 6.2 mm width.

## **4 CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Etoricoxib Tablets is indicated in adults and adolescents 16 years of age and older for the symptomatic relief of osteoarthritis (OA), rheumatoid arthritis (RA), ankylosing spondylitis, and the pain and signs of inflammation associated with acute gouty arthritis.

Etoricoxib Tablets is indicated in adults and adolescents 16 years of age and older for the short-term treatment of moderate pain associated with dental surgery.

The decision to prescribe a selective COX-2 inhibitor should be based on an assessment of the individual patient's overall risks (see sections 4.3, 4.4).

### **4.2 Posology and method of administration**

#### Posology

As the cardiovascular risks of etoricoxib may increase with dose and duration of exposure, the shortest duration possible and the lowest effective daily dose should be used. The patient's need for symptomatic relief and response to therapy should be re-evaluated periodically, especially in patients with osteoarthritis (see sections 4.3, 4.4, 4.8 and 5.1).

#### *Osteoarthritis*

The recommended dose is 30 mg once daily. In some patients with insufficient relief from symptoms, an increased dose of 60 mg once daily may increase efficacy. In the absence of an increase in therapeutic benefit, other therapeutic options should be considered.

#### *Rheumatoid arthritis*

The recommended dose is 60 mg once daily. In some patients with insufficient relief from symptoms, an increased dose of 90 mg once daily may increase efficacy. Once the patient is clinically stabilised, down-titration to a 60 mg once daily dose may be appropriate. In the absence of an increase in therapeutic benefit, other therapeutic options should be considered.

#### *Ankylosing spondylitis*

The recommended dose is 60 mg once daily. In some patients with insufficient relief from symptoms, an increased dose of 90 mg once daily may increase efficacy. Once the patient is clinically stabilised, down-titration to a 60 mg once daily dose may be appropriate. In the absence of an increase in therapeutic benefit, other therapeutic options should be considered.

#### *Acute pain conditions*

For acute pain conditions, Etoricoxib Tablets should be used only for the acute symptomatic period.

#### *Acute gouty arthritis*

The recommended dose is 120 mg once daily. In clinical trials for acute gouty arthritis, Etoricoxib Tablets was given for 8 days.

#### *Postoperative dental surgery pain*

The recommended dose is 90 mg once daily, limited to a maximum of 3 days. Some patients may require other postoperative analgesia in addition to Etoricoxib Tablets during the three day treatment period.

Doses greater than those recommended for each indication have either not demonstrated additional efficacy or have not been studied. Therefore:

The dose for OA should not exceed 60 mg daily.

The dose for RA and ankylosing spondylitis should not exceed 90 mg daily.

The dose for acute gout should not exceed 120 mg daily, limited to a maximum of 8 days treatment.

The dose for postoperative acute dental surgery pain should not exceed 90 mg daily, limited to a maximum of 3 days.

### Special populations

#### *Elderly patients*

No dosage adjustment is necessary for elderly patients. As with other drugs, caution should be exercised in elderly patients (see section 4.4).

#### *Patients with hepatic impairment*

Regardless of indication, in patients with mild hepatic dysfunction (Child-Pugh score 5-6) a dose of 60 mg once daily should not be exceeded. In patients with moderate hepatic dysfunction (Child-Pugh score 7-9), regardless of indication, the dose of 30 mg *once daily* should not be exceeded.

Clinical experience is limited particularly in patients with moderate hepatic dysfunction and caution is advised. There is no clinical experience in patients with severe hepatic dysfunction (Child-Pugh score  $\geq 10$ ); therefore, its use is contraindicated in these patients (see sections 4.3, 4.4 and 5.2).

#### *Patients with renal impairment*

No dosage adjustment is necessary for patients with creatinine clearance  $\geq 30$  ml/min (see section 5.2). The use of Etoricoxib Tablets in patients with creatinine clearance  $< 30$  ml/min is contra-indicated (see sections 4.3 and 4.4).

#### *Paediatric population*

Etoricoxib Tablets is contra-indicated in children and adolescents under 16 years of age (see section 4.3).

#### Method of administration

Etoricoxib Tablets is administered orally and may be taken with or without food. The onset of the effect of the medicinal product may be faster when Etoricoxib Tablets is administered without food. This should be considered when rapid symptomatic relief is needed.

### **4.3 Contraindications**

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Active peptic ulceration or active gastro-intestinal (GI) bleeding.
- Patients who, after taking acetylsalicylic acid or NSAIDs including COX-2 (cyclooxygenase-2) inhibitors, experience bronchospasm, acute rhinitis, nasal polyps, angioneurotic oedema, urticaria, or allergic-type reactions.
- Pregnancy and lactation (see sections 4.6 and 5.3).
- Severe hepatic dysfunction (serum albumin  $< 25$  g/l or Child-Pugh score  $\geq 10$ ).
- Estimated renal creatinine clearance  $< 30$  ml/min.
- Children and adolescents under 16 years of age.
- Inflammatory bowel disease.
- Congestive heart failure (NYHA II-IV).
- Patients with hypertension whose blood pressure is persistently elevated above 140/90 mmHg and has not been adequately controlled.
- Established ischaemic heart disease, peripheral arterial disease, and/or cerebrovascular disease.

## 4.4 Special warnings and precautions for use

### *Gastrointestinal effects*

Upper gastrointestinal complications [perforations, ulcers or bleedings (PUBs)], some of them resulting in fatal outcome, have occurred in patients treated with etoricoxib tablets.

Caution is advised with treatment of patients most at risk of developing a gastrointestinal complication with NSAIDs; the elderly, patients using any other NSAID or acetylsalicylic acid concomitantly or patients with a prior history of gastrointestinal disease, such as ulceration and GI bleeding.

There is a further increase in the risk of gastrointestinal adverse effects (gastrointestinal ulceration or other gastrointestinal complications) when Etoricoxib Tablets is taken concomitantly with acetylsalicylic acid (even at low doses). A significant difference in GI safety between selective COX-2 inhibitors + acetylsalicylic acid vs. NSAIDs + acetylsalicylic acid has not been demonstrated in long-term clinical trials (see section 5.1).

### *Cardiovascular effects*

Clinical trials suggest that the selective COX-2 inhibitor class of drugs may be associated with a risk of thrombotic events (especially myocardial infarction (MI) and stroke), relative to placebo and some NSAIDs. As the cardiovascular risks of Etoricoxib Tablets may increase with dose and duration of exposure, the shortest duration possible and the lowest effective daily dose should be used. The patient's need for symptomatic relief and response to therapy should be re-evaluated periodically, especially in patients with osteoarthritis (see sections 4.2, 4.3, 4.8 and 5.1).

Patients with significant risk factors for cardiovascular events (e.g. hypertension, hyperlipidaemia, diabetes mellitus, smoking) should only be treated with Etoricoxib Tablets after careful consideration (see section 5.1).

COX-2 selective inhibitors are not a substitute for acetylsalicylic acid for prophylaxis of cardiovascular thrombo-embolic diseases because of their lack of antiplatelet effect. Therefore antiplatelet therapies should not be discontinued (see sections above, 4.5 and 5.1.).

### *Renal effects*

Renal prostaglandins may play a compensatory role in the maintenance of renal perfusion. Therefore, under conditions of compromised renal perfusion, administration of Etoricoxib Tablets may cause a reduction in prostaglandin formation and, secondarily, in renal blood flow, and thereby impair renal function. Patients at greatest risk of this response are those with pre-existing significantly impaired renal function, uncompensated heart failure, or cirrhosis. Monitoring of renal function in such patients should be considered.

### *Fluid retention, oedema and hypertension*

As with other medicinal products known to inhibit prostaglandin synthesis, fluid retention, oedema and hypertension have been observed in patients taking etoricoxib. All Nonsteroidal Anti-inflammatory Drugs (NSAIDs), including Etoricoxib Tablets, can be associated with new onset or recurrent congestive heart failure. For

information regarding a dose related response for Etoricoxib Tablets see section 5.1. Caution should be exercised in patients with a history of cardiac failure, left ventricular dysfunction, or hypertension and in patients with pre-existing oedema from any other reason. If there is clinical evidence of deterioration in the condition of these patients, appropriate measures including discontinuation of etoricoxib should be taken.

Etoricoxib Tablets may be associated with more frequent and severe hypertension than some other NSAIDs and selective COX-2 inhibitors, particularly at high doses. Therefore, hypertension should be controlled before treatment with Etoricoxib Tablets (see section 4.3) and special attention should be paid to blood pressure monitoring during treatment with Etoricoxib Tablets. Blood pressure should be monitored within two weeks after initiation of treatment and periodically thereafter. If blood pressure rises significantly, alternative treatment should be considered.

#### *Hepatic effects*

Elevations of alanine aminotransferase (ALT) and/or aspartate aminotransferase (AST) (approximately three or more times the upper limit of normal) have been reported in approximately 1% of patients in clinical trials treated for up to one year with Etoricoxib Tablets 30, 60 and 90 mg daily.

Any patients with symptoms and/or signs suggesting liver dysfunction, or in whom an abnormal liver function test has occurred, should be monitored. If signs of hepatic insufficiency occur, or if persistently abnormal liver function tests (three times the upper limit of normal) are detected, Etoricoxib Tablets should be discontinued.

#### *General*

If during treatment, patients deteriorate in any of the organ system functions described above, appropriate measures should be taken and discontinuation of Etoricoxib Tablets therapy should be considered. Medically appropriate supervision should be maintained when using Etoricoxib Tablets in the elderly and in patients with renal, hepatic, or cardiac dysfunction.

Caution should be used when initiating treatment with Etoricoxib Tablets in patients with dehydration. It is advisable to rehydrate patients prior to starting therapy with Etoricoxib Tablets.

Serious skin reactions, some of them fatal, including exfoliative dermatitis, Stevens-Johnson syndrome, and toxic epidermal necrolysis, have been reported very rarely in association with the use of NSAIDs and some selective COX-2 inhibitors during post-marketing surveillance (see section 4.8). Patients appear to be at highest risk for these reactions early in the course of therapy with the onset of the reaction occurring in the majority of cases within the first month of treatment. Serious hypersensitivity reactions (such as anaphylaxis and angioedema) have been reported in patients receiving Etoricoxib Tablets (see section 4.8). Some selective COX-2 inhibitors have been associated with an increased risk of skin reactions in patients with a history of any drug allergy. Etoricoxib Tablets should be discontinued at the first appearance of skin rash, mucosal lesions, or any other sign of hypersensitivity.

Etoricoxib Tablets may mask fever and other signs of inflammation.

Caution should be exercised when co-administering etoricoxib tablets with warfarin or other oral anticoagulants (see section 4.5).

The use of Etoricoxib Tablets, as with any medicinal product known to inhibit cyclooxygenase / prostaglandin synthesis, is not recommended in women attempting to conceive (see sections 4.6, 5.1, and 5.3).

Etoricoxib Tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

## **4.5 Interaction with other medicinal products and other forms of interaction**

### *Pharmacodynamic interactions*

**Oral anticoagulants:** In subjects stabilised on chronic warfarin therapy, the administration of Etoricoxib Tablets 120 mg daily was associated with an approximate 13% increase in prothrombin time International Normalised Ratio (INR). Therefore, patients receiving oral anticoagulants should be closely monitored for their prothrombin time INR, particularly in the first few days when therapy with Etoricoxib Tablets is initiated or the dose of Etoricoxib Tablets is changed (see section 4.4).

**Diuretics, ACE inhibitors and Angiotensin II Antagonists:** NSAIDs may reduce the effect of diuretics and other antihypertensive drugs. In some patients with compromised renal function (e.g. dehydrated patients or elderly patients with compromised renal function) the co-administration of an ACE inhibitor or Angiotensin II antagonist and agents that inhibit cyclo-oxygenase may result in further deterioration of renal function, including possible acute renal failure, which is usually reversible. These interactions should be considered in patients taking Etoricoxib Tablets concomitantly with ACE inhibitors or angiotensin II antagonists. Therefore, the combination should be administered with caution, especially in the elderly. Patients should be adequately hydrated and consideration should be given to monitoring of renal function after initiation of concomitant therapy, and periodically thereafter.

**Acetylsalicylic Acid:** In a study in healthy subjects, at steady state, Etoricoxib Tablets 120 mg once daily had no effect on the anti-platelet activity of acetylsalicylic acid (81 mg once daily). Etoricoxib Tablets can be used concomitantly with acetylsalicylic acid at doses used for cardiovascular prophylaxis (low-dose acetylsalicylic acid). However, concomitant administration of low-dose acetylsalicylic acid with Etoricoxib Tablets may result in an increased rate of GI ulceration or other complications compared to use of Etoricoxib Tablets. Concomitant administration of Etoricoxib Tablets with doses of acetylsalicylic acid above those for cardiovascular prophylaxis or with other NSAIDs is not recommended (see sections 5.1 and 4.4.).

**Cyclosporin and tacrolimus:** Although this interaction has not been studied with Etoricoxib Tablets, coadministration of cyclosporin or tacrolimus with any NSAID may increase the nephrotoxic effect of cyclosporin or tacrolimus. Renal function should be monitored when Etoricoxib Tablets and either of these drugs is used in combination.

### *Pharmacokinetic interactions*

#### *The effect of etoricoxib tablets on the pharmacokinetics of other drugs*

**Lithium:** NSAIDs decrease lithium renal excretion and therefore increase lithium plasma levels. If necessary, monitor blood lithium closely and adjust the lithium dosage while the combination is being taken and when the NSAID is withdrawn.

*Methotrexate:* Two studies investigated the effects of Etoricoxib Tablets 60, 90 or 120 mg administered once daily for seven days in patients receiving once-weekly methotrexate doses of 7.5 to 20 mg for rheumatoid arthritis. Etoricoxib Tablets at 60 and 90 mg had no effect on methotrexate plasma concentrations or renal clearance. In one study, Etoricoxib Tablets 120 mg had no effect, but in the other study, Etoricoxib Tablets 120 mg increased methotrexate plasma concentrations by 28% and reduced renal clearance of methotrexate by 13%. Adequate monitoring for methotrexate-related toxicity is recommended when Etoricoxib Tablets and methotrexate are administered concomitantly.

*Oral contraceptives:* Etoricoxib Tablets 60 mg given concomitantly with an oral contraceptive containing 35 micrograms ethinyl estradiol (EE) and 0.5 to 1 mg norethindrone for 21 days increased the steady state AUC<sub>0-24hr</sub> of EE by 37%. Etoricoxib Tablets 120 mg given with the same oral contraceptive concomitantly or separated by 12 hours, increased the steady state AUC<sub>0-24hr</sub> of EE by 50 to 60%. This increase in EE concentration should be considered when selecting an oral contraceptive for use with etoricoxib. An increase in EE exposure can increase the incidence of adverse events associated with oral contraceptives (e.g., venous thrombo-embolic events in women at risk).

*Hormone Replacement Therapy (HRT):* Administration of Etoricoxib Tablets 120 mg with hormone replacement therapy consisting of conjugated estrogens (0.625 mg PREMARIN<sup>TM</sup>) for 28 days, increased the mean steady state AUC<sub>0-24hr</sub> of unconjugated estrone (41%), equilin (76%), and 17- $\beta$ -estradiol (22%). The effect of the recommended chronic doses of Etoricoxib Tablets (30, 60, and 90 mg) has not been studied. The effects of Etoricoxib Tablets 120 mg on the exposure (AUC<sub>0-24hr</sub>) to these estrogenic components of PREMARIN were less than half of those observed when PREMARIN was administered alone and the dose was increased from 0.625 to 1.25 mg. The clinical significance of these increases is unknown, and higher doses of PREMARIN were not studied in combination with etoricoxib. These increases in estrogenic concentration should be taken into consideration when selecting post-menopausal hormone therapy for use with Etoricoxib Tablets because the increase in oestrogen exposure might increase the risk of adverse events associated with HRT.

*Prednisone/prednisolone:* In drug-interaction studies, Etoricoxib Tablets did not have clinically important effects on the pharmacokinetics of prednisone/prednisolone.

*Digoxin:* Etoricoxib tablets 120 mg administered once daily for 10 days to healthy volunteers did not alter the steady-state plasma AUC<sub>0-24hr</sub> or renal elimination of digoxin. There was an increase in digoxin C<sub>max</sub> (approximately 33%). This increase is not generally important for most patients. However, patients at high risk of digoxin toxicity should be monitored for this when Etoricoxib Tablets and digoxin are administered concomitantly.

#### *Effect of Etoricoxib Tablets on drugs metabolised by sulfotransferases*

Etoricoxib Tablets is an inhibitor of human sulfotransferase activity, particularly SULT1E1, and has been shown to increase the serum concentrations of ethinyl estradiol. While knowledge about effects of multiple sulfotransferases is presently limited and the clinical consequences for many drugs are still being examined, it may be prudent to exercise care when administering Etoricoxib Tablets concurrently with other drugs primarily metabolised by human sulfotransferases (e.g., oral salbutamol and minoxidil).

#### *Effect of Etoricoxib Tablets on drugs metabolised by CYP isoenzymes*

Based on *in vitro* studies, Etoricoxib Tablets is not expected to inhibit cytochromes P450 (CYP) 1A2, 2C9, 2C19, 2D6, 2E1 or 3A4. In a study in healthy subjects, daily administration of Etoricoxib Tablets 120 mg did not alter hepatic CYP3A4 activity as assessed by the erythromycin breath test.

#### *Effects of other drugs on the pharmacokinetics of Etoricoxib Tablets*

The main pathway of Etoricoxib Tablets metabolism is dependent on CYP enzymes. CYP3A4 appears to contribute to the metabolism of etoricoxib *in vivo*. *In vitro* studies indicate that CYP2D6, CYP2C9, CYP1A2 and CYP2C19 also can catalyse the main metabolic pathway, but their quantitative roles have not been studied *in vivo*.

*Ketoconazole*: Ketoconazole, a potent inhibitor of CYP3A4, dosed at 400 mg once a day for 11 days to healthy volunteers, did not have any clinically important effect on the single-dose pharmacokinetics of 60 mg Etoricoxib Tablets (43% increase in AUC).

*Voriconazole and Miconazole*: Co-administration of either oral voriconazole or topical miconazole oral gel, strong CYP3A4 inhibitors, with Etoricoxib Tablets caused a slight increase in exposure to etoricoxib, but is not considered to be clinically meaningful based on published data.

*Rifampicin*: Co-administration of Etoricoxib Tablets with rifampicin, a potent inducer of CYP enzymes, produced a 65% decrease in Etoricoxib Tablets plasma concentrations. This interaction may result in recurrence of symptoms when Etoricoxib Tablets is co-administered with rifampicin. While this information may suggest an increase in dose, doses of etoricoxib greater than those listed for each indication have not been studied in combination with rifampicin and are therefore not recommended (see section 4.2).

*Antacids*: Antacids do not affect the pharmacokinetics of Etoricoxib Tablets to a clinically relevant extent.

## **4.6 Fertility, pregnancy and lactation**

### *Pregnancy*

No clinical data on exposed pregnancies are available for Etoricoxib Tablets. Studies in animals have shown reproductive toxicity (see section 5.3). The potential for human risk in pregnancy is unknown. Etoricoxib Tablets, as with other medicinal products inhibiting prostaglandin synthesis, may cause uterine inertia and premature closure of the ductus arteriosus during the last trimester. Etoricoxib Tablets is contraindicated in pregnancy (see section 4.3). If a woman becomes pregnant during treatment, etoricoxib must be discontinued.

### *Breastfeeding*

It is not known whether Etoricoxib Tablets is excreted in human milk. Etoricoxib Tablets is excreted in the milk of lactating rats. Women who use etoricoxib must not breast feed (see sections 4.3 and 5.3).

### *Fertility*

The use of Etoricoxib Tablets, as with any drug substance known to inhibit COX-2, is not recommended in women attempting to conceive.

#### 4.7 Effects on ability to drive and use machines

Patients who experience dizziness, vertigo or somnolence while taking Etoricoxib Tablets should refrain from driving or operating machinery.

#### 4.8 Undesirable effects

##### *Summary of the safety profile*

In clinical trials, Etoricoxib Tablets was evaluated for safety in 9,295 individuals, including 6,757 patients with OA, RA, chronic low back pain or ankylosing spondylitis (approximately 600 patients with OA or RA were treated for one year or longer).

In clinical studies, the undesirable effects profile was similar in patients with OA or RA treated with Etoricoxib Tablets for one year or longer.

In a clinical study for acute gouty arthritis, patients were treated with Etoricoxib Tablets 120 mg once daily for eight days. The adverse experience profile in this study was generally similar to that reported in the combined OA, RA, and chronic low back pain studies.

In a cardiovascular safety outcomes programme of pooled data from three active comparator controlled trials, 17,412 patients with OA or RA were treated with Etoricoxib Tablets (60 mg or 90 mg) for a mean duration of approximately 18 months. The safety data and details from this programme are presented in section 5.1.

In clinical studies for acute postoperative dental pain following surgery including 614 patients treated with Etoricoxib Tablets (90 mg or 120 mg), the adverse experience profile in these studies was generally similar to that reported in the combined OA, RA, and chronic low back pain studies.

##### *Tabulated list of adverse reactions*

The following undesirable effects were reported at an incidence greater than placebo in clinical trials in patients with OA, RA, chronic low back pain or ankylosing spondylitis treated with Etoricoxib Tablets 30 mg, 60 mg or 90 mg up to the recommended dose for up to 12 weeks; in the MEDAL Programme studies for up to 3½ years; in short term acute pain studies for up to 7 days; or in post-marketing experience (see Table 1):

**Table 1:**

<b>System Organ Class</b>	<b>Adverse Reactions</b>	<b>Frequency Category*</b>
<b><i>Infections and infestations</i></b>	alveolar osteitis	Common
	gastroenteritis, upper respiratory infection, urinary tract infection	Uncommon
<b><i>Blood and lymphatic system disorders</i></b>	anaemia (primarily associated with gastrointestinal bleeding), leukopenia, thrombocytopenia	Uncommon
<b><i>Immune system disorders</i></b>	hypersensitivity <sup>*B</sup>	Uncommon
	angioedema/anaphylactic /anaphylactoid	Rare

	reactions including shock <sup>‡</sup>	
<b>Metabolism and nutrition disorders</b>	oedema/fluid retention	Common
	appetite increase or decrease, weight gain	Uncommon
<b>Psychiatric disorders</b>	anxiety, depression, mental acuity decreased, hallucinations <sup>‡</sup>	Uncommon
	confusion <sup>‡</sup> , restlessness <sup>‡</sup>	Rare
<b>Nervous system disorders</b>	dizziness, headache	Common
	dysgeusia, insomnia, paresthaesia/hypaesthesia, somnolence	Uncommon
<b>Eye disorders</b>	blurred vision, conjunctivitis	Uncommon
<b>Ear and labyrinth disorders</b>	tinnitus, vertigo	Uncommon
<b>Cardiac disorders</b>	palpitations, arrhythmia <sup>‡</sup>	Common
	atrial fibrillation, tachycardia <sup>‡</sup> , congestive heart failure, non-specific ECG changes, angina pectoris <sup>‡</sup> , myocardial infarction <sup>§</sup>	Uncommon
<b>Vascular disorders</b>	Hypertension	Common
	flushing, cerebrovascular accident <sup>§</sup> , transient ischaemic attack, hypertensive crisis <sup>‡</sup> , vasculitis <sup>‡</sup>	Uncommon
<b>Respiratory, thoracic and mediastinal disorders</b>	bronchospasm <sup>‡</sup>	Common
	cough, dyspnoea, epistaxis	Uncommon
<b>Gastrointestinal disorders</b>	abdominal pain	Very common
	Constipation, flatulence, gastritis, heartburn/acid reflux, diarrhea, dyspepsia/epigastric discomfort, nausea, vomiting, oesophagitis, oral ulcer	Common
	abdominal distention, bowel movement pattern change, dry mouth, gastroduodenal ulcer, peptic ulcers including gastrointestinal perforation and bleeding, irritable bowel syndrome, pancreatitis <sup>‡</sup>	Uncommon
<b>Hepatobiliary disorders</b>	ALT increased, AST increased	Common
	hepatitis <sup>‡</sup>	Rare
	hepatic failure <sup>‡</sup> , jaundice <sup>‡</sup>	Rare <sup>†</sup>
<b>Skin and subcutaneous tissue disorders</b>	ecchymosis	Common
	facial oedema, pruritus, rash, erythema <sup>‡</sup> , urticaria <sup>‡</sup>	Uncommon
	Stevens-Johnson syndrome <sup>‡</sup> , toxic epidermal necrolysis <sup>‡</sup> , fixed drug eruption <sup>‡</sup>	Rare <sup>†</sup>
<b>Musculoskeletal and connective tissue disorders</b>	muscular cramp/spasm, musculoskeletal pain/stiffness	Uncommon
<b>Renal and urinary disorders</b>	proteinuria, serum creatinine increased, renal failure/renal insufficiency <sup>‡</sup> (see section 4.4)	Uncommon
<b>General disorders and administration site conditions</b>	asthenia/fatigue, flu-like disease	Common
	chest pain	Uncommon
<b>Investigations</b>	blood urea nitrogen increased, creatine	Uncommon

	phosphokinase increased, hyperkalaemia, uric acid increased	
	blood sodium decreased	Rare

\*Frequency Category: Defined for each Adverse Experience Term by the incidence reported in the clinical trials data base: Very Common ( $\geq 1/10$ ), Common ( $\geq 1/100$  to  $< 1/10$ ), Uncommon ( $\geq 1/1000$  to  $< 1/100$ ), Rare ( $\geq 1/10,000$  to  $< 1/1000$ ), Very Rare ( $< 1/10,000$ ).

‡ This adverse reaction was identified through post-marketing surveillance. Its reported frequency has been estimated based upon the highest frequency observed across clinical trial data pooled by indication and approved dose.

†The frequency category of “Rare” was defined per the Summary of Product Characteristics (SmPC) guidance (rev.2, Sept 2009) on the basis of an estimated upper bound of the 95% confidence interval for 0 events given the number of subjects treated with Etoricoxib Tablets in the analysis of the Phase III data pooled by dose and indication (n=15,470).

<sup>b</sup> Hypersensitivity includes the terms "allergy", "drug allergy", "drug hypersensitivity", "hypersensitivity", "hypersensitivity NOS", "hypersensitivity reaction" and nonspecific allergy".

<sup>§</sup>Based on analyses of long-term placebo and active controlled clinical trials, selective COX-2 inhibitors have been associated with an increased risk of serious thrombotic arterial events, including myocardial infarction and stroke. The absolute risk increase for such events is unlikely to exceed 1% per year based on existing data (uncommon).

The following serious undesirable effects have been reported in association with the use of NSAIDs and cannot be ruled out for Etoricoxib Tablets: nephrotoxicity including interstitial nephritis and nephrotic syndrome.

#### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard).

## **4.9 Overdose**

In clinical studies, administration of single doses of Etoricoxib Tablets up to 500 mg and multiple doses up to 150 mg/day for 21 days did not result in significant toxicity. There have been reports of acute overdosage with Etoricoxib Tablets, although adverse experiences were not reported in the majority of cases. The most frequently observed adverse experiences were consistent with the safety profile for Etoricoxib Tablets (e.g. gastrointestinal events, cardiorenal events).

In the event of overdose, it is reasonable to employ the usual supportive measures, e.g., remove unabsorbed material from the GI tract, employ clinical monitoring, and institute supportive therapy, if required.

Etoricoxib Tablets is not dialysable by haemodialysis; it is not known whether Etoricoxib Tablets is dialysable by peritoneal dialysis.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

*Pharmacotherapeutic group: Anti-inflammatory and antirheumatic products, non-steroids, coxibs, ATC code: M01AH05*

#### Mechanism of Action

Etoricoxib is an oral, selective cyclo-oxygenase-2 (COX-2) inhibitor within the clinical dose range.

Across clinical pharmacology studies, Etoricoxib produced dose-dependent inhibition of COX-2 without inhibition of COX-1 at doses up to 150 mg daily. Etoricoxib did not inhibit gastric prostaglandin synthesis and had no effect on platelet function.

Cyclooxygenase is responsible for generation of prostaglandins. Two isoforms, COX-1 and COX-2, have been identified. COX-2 is the isoform of the enzyme that has been shown to be induced by pro-inflammatory stimuli and has been postulated to be primarily responsible for the synthesis of prostanoid mediators of pain, inflammation, and fever. COX-2 is also involved in ovulation, implantation and closure of the ductus arteriosus, regulation of renal function, and central nervous system functions (fever induction, pain perception and cognitive function). It may also play a role in ulcer healing. COX-2 has been identified in tissue around gastric ulcers in man but its relevance to ulcer healing has not been established.

#### Clinical efficacy and safety

##### Efficacy

In patients with osteoarthritis (OA), Etoricoxib Tablets 60 mg once daily provided significant improvements in pain and patient assessments of disease status. These beneficial effects were observed as early as the second day of therapy and maintained for up to 52 weeks. Studies with Etoricoxib Tablets 30 mg once daily demonstrated efficacy superior to placebo over a 12 week treatment period (using similar assessments as the above studies). In a dose ranging study, Etoricoxib Tablets 60 mg demonstrated significantly greater improvement than 30 mg for all 3 primary endpoints over 6 weeks of treatment. The 30 mg dose has not been studied in osteoarthritis of hands.

In patients with rheumatoid arthritis (RA), Etoricoxib Tablets 60 mg and 90 mg once daily both provided significant improvements in pain, inflammation, and mobility. In studies evaluating the 60 mg and 90 mg dose, these beneficial effects were maintained over the 12-week treatment periods. In a study evaluating the 60 mg dose compared to the 90 mg dose, Etoricoxib Tablets 60 mg once daily and 90 mg once daily were both more effective than placebo. The 90 mg dose was superior to the 60 mg dose for Patient Global Assessment of Pain (0-100mm visual analogue scale), with an average improvement of -2.71 mm (95% CI: -4.98 mm, -0.45 mm).

In patients experiencing attacks of acute gouty arthritis, Etoricoxib Tablets 120 mg once daily over an eight-day treatment period, relieved moderate to extreme joint pain

and inflammation comparable to indomethacin 50 mg three times daily. Pain relief was observed as early as four hours after initiation of treatment.

In patients with ankylosing spondylitis, etoricoxib tablets 90 mg once daily provided significant improvements in spine pain, inflammation, stiffness and function. The clinical benefit of etoricoxib was observed as early as the second day of therapy after initiation of treatment and was maintained throughout the 52-week treatment period. In a second study evaluating the 60 mg dose compared to the 90 mg dose, Etoricoxib Tablets 60 mg daily and 90 mg daily demonstrated similar efficacy compared to naproxen 1,000 mg daily. Among inadequate responders to 60 mg daily for 6 weeks, dose escalation to 90 mg daily improved spinal pain intensity score (0-100 mm visual analogue scale) compared to continuing on 60 mg daily, with an average improvement of -2.70 mm (95% CI: -4.88 mm, -0.52 mm).

In a clinical study evaluating postoperative dental pain, Etoricoxib Tablets 90 mg was administered once daily for up to three days. In the subgroup of patients with moderate pain at baseline, Etoricoxib Tablets 90 mg demonstrated a similar analgesic effect to that of ibuprofen 600 mg (16.11 vs. 16.39; P=0.722), and greater than that of paracetamol/codeine 600 mg/60 mg (11.00; P<0.001) and placebo (6.84; P<0.001) as measured by total pain relief over the first 6 hours (TOPAR6). The proportion of patients reporting rescue medication usage within the first 24 hours of dosing was 40.8% for Etoricoxib Tablets 90 mg, 25.5% for ibuprofen 600 mg Q6h, and 46.7% for paracetamol/codeine 600 mg/60 mg Q6h compared to 76.2% for placebo. In this study, the median onset of action (perceptible pain relief) of 90 mg Etoricoxib Tablets was 28 minutes after dosing.

## Safety

### Multinational Etoricoxib Tablets and Diclofenac Arthritis Long-term (MEDAL) Programme

The MEDAL Programme was a prospectively designed Cardiovascular (CV) Safety Outcomes Programme of pooled data from three randomized, double-blind active comparator controlled trials, the MEDAL study, EDGE II and EDGE.

The MEDAL Study, was an endpoint driven CV Outcomes study in 17,804 OA and 5,700 RA patients treated with Etoricoxib Tablets 60 (OA) or 90 mg (OA and RA) or diclofenac 150 mg daily for a mean period of 20.3 months (maximum of 42.3 months, median 21.3 months). In this trial, only serious adverse events and discontinuations due to any adverse events were recorded.

The EDGE and EDGE II studies compared the gastrointestinal tolerability of Etoricoxib Tablets versus diclofenac. The EDGE study included 7,111 OA patients treated with a dose of Etoricoxib Tablets 90 mg daily (1.5 times the dose recommended for OA) or diclofenac 150 mg daily for a mean period of 9.1 months (maximum 16.6 months, median 11.4 months). The EDGE II study included 4,086 RA patients treated with etoricoxib 90 mg daily or diclofenac 150 mg daily for a mean period of 19.2 months (maximum 33.1 months, median 24 months).

In the pooled MEDAL Programme, 34,701 patients with OA or RA were treated for a mean duration of 17.9 months (maximum 42.3 months, median 16.3 months) with approximately 12,800 patients receiving treatment for more than 24 months. Patients enrolled in the Programme had a wide range of cardiovascular and gastrointestinal risk factors at baseline. Patients with a recent history of myocardial infarction, coronary artery bypass grafting or percutaneous coronary intervention within 6

months preceding enrollment were excluded. Use of gastroprotective agents and low dose aspirin were permitted in the studies.

**Overall Safety:**

There was no significant difference between Etoricoxib Tablets and diclofenac in the rate of cardiovascular thrombotic events. Cardioresenal adverse events were observed more frequently with Etoricoxib Tablets than with diclofenac, and this effect was dose-dependent (see specific results below). Gastrointestinal and hepatic adverse events were observed significantly more frequently with diclofenac than Etoricoxib Tablets. The incidence of adverse experiences in EDGE and EDGE II and of adverse experiences considered serious or resulting in discontinuation in the MEDAL study was higher with Etoricoxib Tablets than diclofenac.

**Cardiovascular safety results**

The rate of confirmed thrombotic cardiovascular serious adverse events (consisting of cardiac, cerebrovascular, and peripheral vascular events) was comparable between Etoricoxib Tablets and diclofenac, and data are summarized in the table below. There were no statistically significant differences in thrombotic event rates between Etoricoxib Tablets and diclofenac across all subgroups analyzed including patient categories across a range of baseline cardiovascular risk. When considered separately, the relative risks for confirmed thrombotic cardiovascular serious adverse events with Etoricoxib Tablets 60 mg or 90 mg compared with diclofenac 150 mg were similar.

<b>Table 2: Rates of Confirmed Thrombotic CV Events (Pooled MEDAL Programme)</b>			
	<b>Etoricoxib (N=16,819) 25,836 Patient- Years</b>	<b>Diclofenac (N=16,483) 24,766 Patient- Years</b>	<b>Between Treatment Comparison</b>
	<b>Rate<sup>†</sup> (95% CI)</b>	<b>Rate<sup>†</sup> (95% CI)</b>	<b>Relative Risk (95% CI)</b>
<b>Confirmed Thrombotic Cardiovascular Serious Adverse Events</b>			
Per-protocol	1.24 (1.11, 1.38)	1.30 (1.17, 1.45)	0.95 (0.81, 1.11)
Intent-to-treat	1.25 (1.14, 1.36)	1.19 (1.08, 1.30)	1.05 (0.93, 1.19)
<b>Confirmed Cardiac Events</b>			
Per-protocol	0.71 (0.61, 0.82)	0.78 (0.68, 0.90)	0.90 (0.74, 1.10)
Intent-to-treat	0.69 (0.61, 0.78)	0.70 (0.62, 0.79)	0.99 (0.84, 1.17)
<b>Confirmed Cerebrovascular Events</b>			
Per-protocol	0.34 (0.28, 0.42)	0.32 (0.25, 0.40)	1.08 (0.80, 1.46)
Intent-to-treat	0.33 (0.28, 0.39)	0.29 (0.24, 0.35)	1.12 (0.87, 1.44)
<b>Confirmed Peripheral Vascular Events</b>			
Per-protocol	0.20 (0.15, 0.27)	0.22 (0.17, 0.29)	0.92 (0.63, 1.35)
Intent-to-treat	0.24 (0.20, 0.30)	0.23 (0.18, 0.28)	1.08 (0.81, 1.44)
<sup>†</sup> Events per 100 Patient-Years; CI=confidence interval  N=total number of patients included in Per-protocol population  Per-protocol: all events on study therapy or within 14 days of discontinuation (excluded: patients who took < 75% of their study medication or took non-study NSAIDs >10% of the time).  Intent-to-treat: all confirmed events up to the end of the trial (included patients potentially exposed to non-study interventions following discontinuation of study medication). Total			

number of patients randomised, n= 17,412 on Etoricoxib Tablets and 17,289 on diclofenac.

CV mortality, as well as overall mortality, was similar between the Etoricoxib Tablets and diclofenac treatment groups.

#### Cardiorenal Events:

Approximately 50% of patients enrolled in the MEDAL study had a history of hypertension at baseline. In the study, the incidence of discontinuations due to hypertension-related adverse events was statistically significantly higher for Etoricoxib Tablets than for diclofenac. The incidence of congestive heart failure adverse events (discontinuations and serious events) occurred at similar rates on Etoricoxib Tablets 60 mg compared to diclofenac 150 mg but was higher for Etoricoxib Tablets 90 mg compared to diclofenac 150 mg (statistically significant for 90 mg Etoricoxib Tablets vs. 150 mg diclofenac in MEDAL OA cohort). The incidence of confirmed congestive heart failure adverse events (events that were serious and resulted in hospitalisation or a visit to an emergency department) was non-significantly higher with Etoricoxib Tablets than diclofenac 150 mg, and this effect was dose-dependent. The incidence of discontinuations due to oedema-related adverse events was higher for Etoricoxib Tablets than diclofenac 150 mg, and this effect was dose-dependent (statistically significant for Etoricoxib Tablets 90 mg, but not for Etoricoxib Tablets 60 mg).

The cardiorenal results for EDGE and EDGE II were consistent with those described for the MEDAL Study.

In the individual MEDAL Programme studies, for Etoricoxib Tablets (60 mg or 90 mg), the absolute incidence of discontinuation in any treatment group was up to 2.6% for hypertension, up to 1.9% for oedema, and up to 1.1% for congestive heart failure, with higher rates of discontinuation observed with Etoricoxib Tablets 90 mg than Etoricoxib Tablets 60 mg.

#### MEDAL Programme Gastrointestinal Tolerability Results:

A significantly lower rate of discontinuations of treatment for any clinical (e.g., dyspepsia, abdominal pain, ulcer) GI adverse event was observed with etoricoxib compared with diclofenac within each of the three component studies of the MEDAL Programme. The rates of discontinuations due to adverse clinical GI events per hundred patient-years over the entire period of study were as follows: 3.23 for Etoricoxib Tablets and 4.96 for diclofenac in the MEDAL Study; 9.12 with Etoricoxib Tablets and 12.28 with diclofenac in the EDGE study; and 3.71 with Etoricoxib Tablets and 4.81 with diclofenac in the EDGE II study.

#### MEDAL Programme Gastrointestinal Safety Results:

Overall upper GI events were defined as perforations, ulcers and bleeds. The subset of overall upper GI events considered complicated included perforations, obstructions, and complicated bleeding; the subset of upper GI events considered uncomplicated included uncomplicated bleeds and uncomplicated ulcers. A significantly lower rate of overall upper GI events was observed with etoricoxib compared to diclofenac. There was no significant difference between Etoricoxib Tablets and diclofenac in the rate of complicated events. For the subset of upper GI haemorrhage events (complicated and uncomplicated combined), there was no significant difference between Etoricoxib Tablets and diclofenac. The upper GI benefit for etoricoxib compared with diclofenac was not statistically significant in patients taking concomitant low-dose aspirin (approximately 33% of patients).

The rates per hundred patient-years of confirmed complicated and uncomplicated upper GI clinical events (perforations, ulcers and bleeds (PUBs)) were 0.67 (95% CI 0.57, 0.77) with Etoricoxib Tablets and 0.97 (95% CI 0.85, 1.10) with diclofenac, yielding a relative risk of 0.69 (95% CI 0.57, 0.83).

The rate for confirmed upper GI events in elderly patients was evaluated and the largest reduction was observed in patients  $\geq 75$  years of age (1.35 [95% CI 0.94, 1.87] vs. 2.78 [95% CI 2.14, 3.56] events per hundred patient-years for etoricoxib and diclofenac, respectively).

The rates of confirmed lower GI clinical events (small or large bowel perforation, obstruction, or haemorrhage, (POBs)) were not significantly different between Etoricoxib Tablets and diclofenac.

#### MEDAL Programme Hepatic Safety Results:

Etoricoxib Tablets was associated with a statistically significantly lower rate of discontinuations due to hepatic-related adverse experiences than diclofenac. In the pooled MEDAL Programme, 0.3% of patients on Etoricoxib Tablets and 2.7% of patients on diclofenac discontinued due to hepatic-related adverse experiences. The rate per hundred patient-years was 0.22 on Etoricoxib Tablets and 1.84 for diclofenac (p-value was  $<0.001$  for etoricoxib vs. diclofenac). However, most hepatic adverse experiences in the MEDAL Programme were non-serious.

#### Additional Thrombotic Cardiovascular Safety Data

In clinical studies excluding the MEDAL Programme Studies, approximately 3,100 patients were treated with Etoricoxib Tablets  $\geq 60$  mg daily for 12 weeks or longer. There was no discernible difference in the rate of confirmed serious thrombotic cardiovascular events between patients receiving Etoricoxib Tablets  $\geq 60$  mg, placebo, or non-naproxen NSAIDs. However, the rate of these events was higher in patients receiving Etoricoxib Tablets compared with those receiving naproxen 500 mg twice daily. The difference in antiplatelet activity between some COX-1 inhibiting NSAIDs and selective COX-2 inhibitors may be of clinical significance in patients at risk of thrombo-embolic events. Selective COX-2 inhibitors reduce the formation of systemic (and therefore possibly endothelial) prostacyclin without affecting platelet thromboxane. The clinical relevance of these observations has not been established.

#### Additional Gastrointestinal Safety Data

In two 12-week double-blind endoscopy studies, the cumulative incidence of gastroduodenal ulceration was significantly lower in patients treated with Etoricoxib Tablets 120 mg once daily than in patients treated with either naproxen 500 mg twice

daily or ibuprofen 800 mg three times daily. Etoricoxib had a higher incidence of ulceration as compared to placebo.

#### Renal Function Study in the Elderly

A randomized, double-blind, placebo-controlled, parallel-group study evaluated the effects of 15 days of treatment of Etoricoxib Tablets (90 mg), celecoxib (200 mg bid), naproxen (500 mg bid) and placebo on urinary sodium excretion, blood pressure, and other renal function parameters in subjects 60 to 85 years of age on a 200-mEq/day sodium diet. Etoricoxib Tablets, celecoxib, and naproxen had similar effects on urinary sodium excretion over the 2 weeks of treatment. All active comparators showed an increase relative to placebo with respect to systolic blood pressures; however, Etoricoxib Tablets was associated with a statistically significant increase at Day 14 when compared to celecoxib and naproxen (mean change from baseline for systolic blood pressure: etoricoxib 7.7 mmHg, celecoxib 2.4 mmHg, naproxen 3.6 mmHg).

## 5.2 Pharmacokinetic properties

### *Absorption*

Orally administered Etoricoxib Tablets is well absorbed. The absolute bioavailability is approximately 100%. Following 120 mg once-daily dosing to steady state, the peak plasma concentration (geometric mean  $C_{max}$  = 3.6  $\mu\text{g/ml}$ ) was observed at approximately 1 hour ( $T_{max}$ ) after administration to fasted adults. The geometric mean area under the curve ( $AUC_{0-24hr}$ ) was 37.8  $\mu\text{g}\cdot\text{hr/ml}$ . The pharmacokinetics of Etoricoxib Tablets are linear across the clinical dose range.

Dosing with food (a high-fat meal) had no effect on the extent of absorption of Etoricoxib Tablets after administration of a 120-mg dose. The rate of absorption was affected, resulting in a 36% decrease in  $C_{max}$  and an increase in  $T_{max}$  by 2 hours. These data are not considered clinically significant. In clinical trials, Etoricoxib Tablets was administered without regard to food intake.

### *Distribution*

Etoricoxib Tablets is approximately 92% bound to human plasma protein over the range of concentrations of 0.05 to 5  $\mu\text{g/ml}$ . The volume of distribution at steady state ( $V_{dss}$ ) was approximately 1,20l in humans.

Etoricoxib Tablets crosses the placenta in rats and rabbits, and the blood-brain barrier in rats.

### *Biotransformation*

Etoricoxib Tablets is extensively metabolised with <1% of a dose recovered in urine as the parent drug. The major route of metabolism to form the 6'-hydroxymethyl derivative is catalyzed by CYP enzymes. CYP3A4 appears to contribute to the metabolism of etoricoxib *in vivo*. *In vitro* studies indicate that CYP2D6, CYP2C9, CYP1A2 and CYP2C19 also can catalyse the main metabolic pathway, but their quantitative roles *in vivo* have not been studied.

Five metabolites have been identified in man. The principal metabolite is the 6'-carboxylic acid derivative of Etoricoxib Tablets formed by further oxidation of the 6'-hydroxymethyl derivative. These principal metabolites either demonstrate no

measurable activity or are only weakly active as COX-2 inhibitors. None of these metabolites inhibit COX-1.

#### *Elimination*

Following administration of a single 25-mg radiolabeled intravenous dose of etoricoxib tablets to healthy subjects, 70% of radioactivity was recovered in urine and 20% in faeces, mostly as metabolites. Less than 2% was recovered as unchanged drug.

Elimination of Etoricoxib Tablets occurs almost exclusively through metabolism followed by renal excretion. Steady state concentrations Etoricoxib Tablets are reached within seven days of once daily administration of 120 mg, with an accumulation ratio of approximately 2, corresponding to a half-life of approximately 22 hours. The plasma clearance after a 25-mg intravenous dose is estimated to be approximately 50 ml/min.

#### *Characteristics in patients*

*Elderly patients:* Pharmacokinetics in the elderly (65 years of age and older) are similar to those in the young.

*Gender:* The pharmacokinetics of Etoricoxib Tablets are similar between men and women.

*Hepatic impairment:* Patients with mild hepatic dysfunction (Child-Pugh score 5-6) administered Etoricoxib Tablets 60 mg once daily had an approximately 16% higher mean AUC as compared to healthy subjects given the same regimen. Patients with moderate hepatic dysfunction (Child-Pugh score 7-9) administered Etoricoxib Tablets 60 mg **every other day** had similar mean AUC to the healthy subjects given Etoricoxib Tablets 60 mg once daily; Etoricoxib Tablets 30 mg once daily has not been studied in this population. There are no clinical or pharmacokinetic data in patients with severe hepatic dysfunction (Child-Pugh score  $\geq 10$ ). (See sections 4.2 and 4.3.)

*Renal impairment:* The pharmacokinetics of a single dose of Etoricoxib Tablets 120 mg in patients with moderate to severe renal insufficiency and patients with end-stage renal disease on haemodialysis were not significantly different from those in healthy subjects. Haemodialysis contributed negligibly to elimination (dialysis clearance approximately 50 ml/min). (See sections 4.3 and 4.4.)

*Paediatric patients:* The pharmacokinetics of Etoricoxib Tablets in paediatric patients (<12 years old) have not been studied.

In a pharmacokinetic study (n=16) conducted in adolescents (aged 12 to 17) the pharmacokinetics in adolescents weighing 40 to 60 kg given Etoricoxib Tablets 60 mg once daily and adolescents >60 kg given etoricoxib 90 mg once daily were similar to the pharmacokinetics in adults given Etoricoxib Tablets 90 mg once daily. Safety and effectiveness of Etoricoxib Tablets in paediatric patients have not been established (see section 4.2).

### **5.3 Preclinical safety data**

In preclinical studies, etoricoxib has been demonstrated not to be genotoxic. Etoricoxib tablets was not carcinogenic in mice. Rats developed hepatocellular and thyroid follicular cell adenomas at >2-times the daily human dose [90 mg] based on systemic exposure when dosed daily for approximately two years. Hepatocellular and thyroid follicular cell adenomas observed in rats are considered to be a consequence of rat-specific mechanism related to hepatic CYP enzyme induction. Etoricoxib Tablets has not been shown to cause hepatic CYP3A enzyme induction in humans.

In the rat, gastrointestinal toxicity of Etoricoxib Tablets increased with dose and exposure time. In the 14-week toxicity study Etoricoxib Tablets caused gastrointestinal ulcers at exposures greater than those seen in man at the therapeutic dose. In the 53- and 106-week toxicity study, gastrointestinal ulcers were also seen at exposures comparable to those seen in man at the therapeutic dose. In dogs, renal and gastrointestinal abnormalities were seen at high exposures.

Etoricoxib Tablets was not teratogenic in reproductive toxicity studies conducted in rats at 15 mg/kg/day (this represents approximately 1.5 times the daily human dose [90 mg] based on systemic exposure). In rabbits, a treatment related increase in cardiovascular malformations was observed at exposure levels below the clinical exposure at the daily human dose (90 mg). However no treatment-related external or skeletal foetal malformations were observed. In rats and rabbits, there was a dose dependent increase in post implantation loss at exposures greater than or equal to 1.5 times the human exposure (see sections 4.3 and 4.6).

Etoricoxib Tablets is excreted in the milk of lactating rats at concentrations approximately two-fold those in plasma. There was a decrease in pup body weight following exposure of pups to milk from dams administered Etoricoxib Tablets during lactation.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

*Core:*

Microcrystalline cellulose  
Calcium hydrogen phosphate anhydrous  
Croscarmellose sodium  
Lactose monohydrate  
Hydroxypropyl cellulose  
Magnesium stearate

*Tablet coating:*

Hypromellose  
Lactose monohydrate  
Titanium dioxide (E171)  
Triacetin  
indigo carmine aluminium lake (E132)  
Iron oxide yellow (E172).

### **6.2 Incompatibilities**

Not applicable

**6.3 Shelf life**

36 months

**6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions

**6.5 Nature and contents of container**

Etoricoxib Tablets are available in OPA/Alu/PVC – Alu blister.

Pack sizes:

30 mg, 60 mg, 90 mg and 120 mg:

Pack sizes of 2, 5, 7, 10, 14, 20, 28, 30, 49, 50, 84, 98, 100 film-coated tablets.

Not all pack sizes may be marketed.

**6.6 Special precautions for disposal**

No special requirements.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

**7 MARKETING AUTHORISATION HOLDER**

Brown & Burk UK Ltd  
5 Marryat Close  
Hounslow West  
Middlesex  
TW4 5DQ  
United Kingdom.

**8 MARKETING AUTHORISATION NUMBER(S)**

PL 25298/0083

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 30/08/2017

Date of Last renewal: 08/12/2021

**10 DATE OF REVISION OF THE TEXT**

08/12/2021